


FLNPS & CBG - Plants for Sale 2026 - Cultivation & Appearance



PLEASE TAKE ONE
















FLNPS plants are donated by our members, who propagate from locally collected seed (wild or garden) and/or divide plants from their own gardens. Cornell Botanic Gardens also grows from locally collected seed.


No wild-collected plants are offered for sale.


NOTES – reg or "regular" is equivalent to typical garden soil, watered as needed.  = Good for pollinators. Spring **Ephemeral**= E (foliage dies back by midsummer). * =Not strictly local native but regional/naturalized.


Herbaceous Plants, including Ferns















| | |
|---|--|
| Actaea pachypoda, doll's eyes (Poisonous) | white fl, white berry, 20-30"; moist, part sun-shade |
| * <i>Actaea podocarpa</i> , mountain bugbane (Appalachia: PA & south) | 3-6', white, late summer bloom; part shade, reg soil |
| <i>Actaea racemosa</i> , black snakeroot (Poisonous) | 3-6', white, late summer bloom; part shade, reg soil |
|  <i>Agostache foeniculum</i> , anise hyssop (Mid-western US native) | blue/pur; 3-4', aromatic foliage; moist -dry, sun -pt shade |
|  <i>Allium cernuum</i> , nodding onion (mid-western source) | pink, 18"; sun -part sun, moist-dry |
| <i>Allium tricoccum</i> , wild leek, ramp <u>E</u> | white, 12", wide leaves; moist shade |
| <i>Anaphalis margaritacea</i> , pearly everlasting | grey/white foliage, white/yel fl; 1-3'; sun-lt shade, dry, poor soil |
| <i>Anemone canadense</i> Canada anemone | white, 6-30", groundcover; moist-wet, limey soil, pt sun-pt shade |
| <i>Anemone virginiana</i> , thimbleweed | white, 12-30"; dry- reg, sun-shade |
| <i>Antennaria parlini</i> , Parlin's pussytoes | white, fuzzy flowers, grey foliage, 3-6"; dry, shallow soil, well drained, sun-p sun |
| <i>Antennaria neglecta</i> , pussytoes, ladies tobacco | white, fuzzy flowers, grey mat forming, 3-6"; dry, shallow soil, well drained, sun-p sun |
| <i>Antennaria plantaginifolia</i> , plantain leaf pussytoes | white, fuzzy flowers, grey-green mat forming foliage, 3-6"; dry, shallow soil, well drained, sun-p sun |
|  <i>Aquilegia canadensis</i> , wild columbine | yel&red, 10-24"; sun-lt shade, well-drained, acid-neutral soil |
| <i>Asarum canadense</i> , wild ginger | foliage, groundcover, 4-6"; shade, regular-wet |
|  <i>Asclepias incarnata</i> , swamp milkweed | pink, 2-4'; sun-p shade, reg-wet soil |
| <i>Asclepias tuberosa</i> butterfly weed | orange, 1-3'; sun-p shade, well drained |
| <i>Athyrium angustum</i> , lady fern | delicate, deciduous, 1-2'; p-shade-shade, moist-dry |
| <i>Baptisia australis</i> , blue false indigo (PA-IN & south) | blue, 2-4', strong taproot; sun-pt sun, reg-moist |
|  <i>Caltha palustris</i> , marsh marigold | yellow, 10-12"; moist-wet, sun - pt sun |
| <i>Campanula americana</i> , tall bellflower (biennial) | blue, p shade, 3-4'; moist - dry, p shade - sun |
| <i>Campanula rotundifolia</i> , harebell | blue, 6"; <u>well</u> -drained, moist-dry, pt shade |
| <i>Collinsonia canadensis</i> , horsebalm | lt yel, fragrant, 2-5'; p shade, rich soil |
|  * <i>Coreopsis palmata</i> , stiff tickseed (native mid-west) | yellow, 3' spreads aggressively; well drained, dry, pt shade |
| * <i>Coreopsis verticillata</i> , threadleaf coreopsis (native mid-Atl & mid-west) | yellow, 2-3'; sun, dry-reg |
| <i>Cystopteris bulbifera</i> , bulblet fern | lacy, 12" drooping fronds, "bulbs" under leaf; shade, moist-dry, well drained |
| <i>Diplazium pycnocarpon</i> (Homalosorus pycnocarpos) narrow leaved glade fern | clumps, slender fronds 1.5-3.5'; rich forest & shale, lime, shade |
|  * <i>Echinacea purpurea</i> , purple coneflower (midwest) | pink w/or center, 18-24"; sun-pt sun; dry-moist soil |
| <i>Eupatorium perfoliatum</i> , boneset | white, 2-4'; sun-p sun, moist-wet soil |
| <i>Euphorbia corollata</i> , flowering spurge | white, airy 3'; sun-pt sun, dry-reg |
| <i>Eurybia divaricata</i> , white wood aster | white, 1-2'; moist-dry, sun-shade |
| <i>Eurybia macrophylla</i> , large-leaved aster | White (some purple), 1-3'; sun - pt shade, reg - dry, well-drained |
| <i>Euthamia graminifolia</i> , narrow-leaved goldenrod | Yellow, 1-4', fragrant, aggressive; moist, sun |


| | | |
|---|--|--|
|  | <i>Eutrochium maculatum</i> , spotted Joe Pye | pink, 5-8'; sun - p shade, moist-regular soil |
| | <i>Gentiana andrewsii</i> , fringe-tip bottle gentian | blue, 10-12"; pt shade, moist, well drained |
| | <i>Geranium maculatum</i> , wild geranium | pink, 12-18"; sun - lt shade, moist soil (in spring) |
| | <i>Geum rivale</i> , water avens | Reddish purple & yellow/peach, 1-2'; sun-pt sun, moist-wet (bog) |
| | <i>Hibiscus moscheutos</i> , swamp rosemallow | large pink, 3-7', <u>slow</u> to emerge in spring; sun, moist-wet, rich |
| | <i>Hydrophyllum virginianum</i> , Virginia waterleaf | white-purple, 1-2'; shade-pt shade, moist |
| | <i>Impatiens capensis</i> , jewelweed (Annual) | orange spotted; 2-4'; moist-wet; p shade-sun |
| | * <i>Iris cristata</i> , dwarf crested iris (PA to GA, w to east OK) | blue, white, <u>lav</u> ; 4-6" sun-pt shade, reg -moist rich soil |
| | <i>Iris versicolor</i> , blue flag | blue-violet, 3-4'; sun, moist - wet soil |
| | <i>Juncus tenuis</i> , path rush | small clump; spiked top; spreads!; moist-dry; sun-pt sun |
| | * <i>Liatris spicata</i> , spiked blazing star (unknown native status NY, but native in surrounding states) | purple spike, 5'; reg-wet, sun-pt sun |
|  | <i>Lobelia cardinalis</i> , cardinal flower | red, 2-4'; sun-lt shade, moist - wet (Biennial -allow reseed!) |
| | <i>Lobelia siphilitica</i> , great blue lobelia | blue or white, 2-3'; sun-pt sun, reg- wet (Biennial -allow reseed!) |
| | <i>Lysimachia ciliata</i> , fringed loosestrife | yellow, 2-4'; pt sun, moist-wet |
| | <i>Maianthemum canadensis</i> , Canada mayflower | white, 5", groundcover; moist -wet, acidic, shade-pt shade |
| | <i>Maianthemum racemosum</i> ; Solomon's plume | 1-3', white, pink-red berries; moist-reg, part sun - shade |
| | <i>Maianthemum stellatum</i> , starry Solomon's plume | 1-3', white flowers, pink berries w/2 different clones; moist, part sun-shade |
| | <i>Matteuccia struthiopteris</i> , ostrich fern | erect, large fern; prominent fertile frond; moist-dry, shade-pt sun |
| | <i>Mertensia virginica</i> , Virginia bluebells <u>E</u> | blue/pink ephemeral, 14-20"; pt sun - shade; moist in spring |
| | <i>Mimulus ringens</i> , Monkey Flower | bl-violet, 2-4'; moist-wet, sun-p sun |
|  | <i>Monarda didyma</i> , bee balm, Oswego tea | red, 3-4'; sun-lt shade, reg - dry |
| | <i>Monarda fistulosa</i> , wild bergamot | pale violet, 3-4'; sun-lt shade, moist-reg |
| | <i>Oenothera pilosella</i> , prairie sundrops | yellow, 1-2'; sun, reg-wet (Poisonous to livestock) |
| | <i>Onoclea sensibilis</i> , sensitive fern | 18-24", distinctive spore frond; reg-wet, shade-pt shade |
| | <i>Opuntia humifusa/caespitosa</i> , Eastern prickly pear | 6"ht, spreads, yel fl; looks dead in winter; well-drained dry, sun, no competition |
| | <i>Packera aurea</i> , golden ragwort | yellow, low groundcover w/15-18" flower stalk; sun- pt shade, dry-moist |
| | * <i>Parthenium integrifolium</i> , wild quinine (East U.S) | white, 2-4', fragrant; sun, dry-reg, tolerant |
|  | * <i>Penstemon digitalis</i> , foxglove beardtongue (eastern US) | white, 2-4'; moist-dry, sun |
|  | * <i>Penstemon digitalis/calycosus</i> , Foxglove beardtongue (eastern US native) | white (digitalis) or purple (calycosus), 2-4'; moist-dry, sun |
|  | <i>Penstemon hirsutus</i> , hairy beardtongue | pur/white, 2'; reg-dry, sun |
| | <i>Phlox divaricata</i> , woods phlox  | pale blue-white, 8-12", reg-moist; p shade-sun |
| | <i>Phlox subulata</i> , moss/creeping phlox  | white/pink/purple, 4-6"; well drained, rocky, sun-pt shade, hates competition |
|  | <i>Physostegia virginiana</i> , Obedient plant  | lavender-pink, 2-3'; sun-p sun; moist |
| | <i>Podophyllum peltatum</i> , mayapple (poisonous) | 1-2', white fl, green fruit, poisonous if unripe; moist-wet, p sun - shade |
| | <i>Polygonatum biflorum</i> , large Solomon's seal | 3', white axial fl, blue berries; regular-moist, p shade- shade |
| | <i>Polypodium virginianum</i> , rock cap or common polypody | Fern; shade; neutral or limy, rocky/very well drained |
| | <i>Polystichum acrostichoides</i> , Christmas fern | evergreen fern, 12-18"; dry-reg soil, shade-pt shade |
|  | <i>Pycnanthemum muticum</i> , Short-toothed mtn mint (threatened) | white-violet w/silvery bracts; 2-4', rhizomatous; sun-p shade; moist |
|  | <i>Pycnanthemum tenuifolium</i> , narrow leaved mtn mint | White, 2-3'; sun-pt shade, dry-reg soil |
|  | <i>Pycnanthemum virginianum</i> , Virginia mountain mint | white, 2-3', fragrant; sun-pt shade, dry-reg soil |
|  | * <i>Ratibida pinnata</i> , gray headed prairie coneflower (native midwest) | yel, 3-5'; reg-dry, sun-pt sun |
|  | * <i>Rudbeckia fulgida</i> , orange coneflower (SE, Gulf) | yellow, 1-2.5' dry-moist, sun-pt sun |

NOTES – reg or "regular" is equivalent to typical garden soil, watered as needed.  = Good for pollinators
Spring **Ephemeral**= E (foliage dies back by midsummer). * =Not strictly local native but regional/naturalized.

| | | |
|---|--|---|
|  | * <i>Rudbeckia hirta</i> , black eyed Susan (Native East U.S) | yellow, biennial, 2-3' sun, moist-dry soil |
| | <i>Rudbeckia laciniata</i> , cut-leaved coneflower  | yellow, 4-6'; sun, wet - dry soil |
|  | * <i>Ruellia humilis</i> , wild petunia (Central & Mid-Atl US) | lilac, 2-4", prostrate; don't crowd; dry-reg, sun-pt sun |
| | <i>Sanguinaria canadense</i> , bloodroot | white, 6", spring ephemeral; neutral soil, pt shade |
|  | <i>Sedum ternatum</i> , woodland stonecrop | white succulent; neutral soil, pt shade, rocky |
| | <i>Silene regia</i> , royal catchfly; (native mid-west) | red fl, 4-5', hummingbirds; sun-p shade, dry-regular |
| | <i>Sisyrinchium sp.</i> , blue-eyed grass | blue, 10"; moist-reg, sun-lt shade |
| | <i>Sium suave</i> , water parsnip | white umbels, 2-6', not poisonous; sun-pt shade, moist-wet |
|  | <i>Solidago arguta</i> , sharp-toothed goldenrod | yellow fl; 2-4'; pt shade-shade; dry-moist |
| | <i>Solidago bicolor</i> , silverrod, white goldenrod | 2-3', white; sun - pt shade, dry-reg, well drained |
| | <i>Solidago caesia</i> , wreath goldenrod | 1.5-4', yel flower all along stem; dry-moist, sun-shade (takes dry shade!) |
| | <i>Solidago flexicaulis</i> , zig-zag goldenrod | 1-4', yellow clusters; moist-dry woodland, sun -shade |
| | <i>Solidago juncea</i> , early goldenrod | yellow, 2-4', pollinators; sun, dry-reg soil |
| | <i>Solidago nemoralis</i> , old-field goldenrod | 6-24", yellow, largish leaves; dry-reg soil, sun |
| | <i>Solidago patula</i> , swamp/rough-leaved goldenrod | yellow, 4-6', large leaves; moist, sun - pt shade |
| | <i>Solidago ptarmicoides</i> , upland white aster | upland white aster; White, 1-2' sun, dry-reg well drained |
|  | <i>Solidago rigida</i> , stiff goldenrod | yel clusters, 4', red fall lvs; moist-dry, sun-p sun |
| | * <i>Stylophorum diphyllum</i> , wood poppy (mid-Atlantic & mid-west US) | yel fl, 12-18"; spreads; pt-full shade, moist-wet |
|  | <i>Symphyotrichum cordifolium</i> , blue wood aster | white-pale blue, 15-24"; pt sh-pt sun, dry-moist |
| | <i>Symphyotrichum laeve</i> , smooth blue aster | lavender, 1-3'; sun-pt sh, reg-dry, well drained, likes lime |
| | <i>Symphyotrichum lateriflorum</i> , calico aster | white, 1-3'; moist-dry, sun-pt shade |
| | <i>Symphyotrichum novae-angliae</i> , New England aster | pink to violet/purple, 2-6'; moist, sun -part sun |
| | <i>Symphyotrichum pilosum v. pringleii</i> , Pringle's aster | many sm white fl, 2-4' x 4' wide, narrow lf; sun, reg-dry; spreads |
| | <i>Symphyotrichum prenanthoides</i> , zig-zag aster | blue/lavender, 24-30"; moist & p-full sun OR part shade if dry |
| | <i>Symphyotrichum puniceum</i> , purple stemmed aster | violet-blue to purple, 4-6'; sun, med-wet |
|  | <i>Symphyotrichum undulatum</i> , wavy-leaved aster | pale bl-viol, 1-3'; pt sun - pt shade, dry-reg soil |
| | <i>Symphyotrichum urophyllum</i> , arrow leaved aster | white, 2-3'; dry, sun -part sun |
| | <i>Thalictrum dioicum</i> , early meadow rue | yel-grn; 1-3', attractive leaves; pt sun-shade; moist in spring |
| | <i>Thalictrum pubescens</i> , tall meadow rue | white, 2-4'; sun-pt shade, moist |
| | <i>Tiarella cordifolia/stolonifera</i> , foam flower | white, 6-8"; shade-pt shade, dry-moist, well drained |
|  | * <i>Tradescantia ohiensis</i> , Ohio spiderwort (native mid-west, SE) | blue, 3'; dry-moist, part shade |
| | <i>Trillium grandiflorum</i> , white trillium | white, 10-12"; shade-pt shade, rich moist soil |
| | <i>Verbena hastata</i> , blue vervain | blue-purple spire, 2-6'; sun, moist-wet |
|  | <i>Vernonia gigantea</i> & <i>V. novaboracensis</i> , ironweed | deep red-purple; 4-8'; p sun, moist-wet |
| | <i>Veronicastrum virginicum</i> , culver's root  | pinkish white, 3-6'; sun-pt sun, moist |
| | <i>Viola canadensis</i> , Canada violet | white w/purple back, stemmed, 10-16"; rich moist soil, shade |
| | <i>Viola pubescens v. pubescens</i> , downy yellow violet | yellow, stemmed; pt shade, med-wet |
| | <i>Viola sororia</i> , common blue violet | purple flowered; moist -dry soil, pt shade - sun |
| | <i>Viola sororia f. priceana</i> (?), white/confederate violet | white flowered, variable purple center, probable hybrid, stemless; moist-dry soil, pt shade-sun |
|  | <i>Viola striata</i> , creamy violet | white, part shade, 8-12"; moist soil, shade - pt shade |
| | <i>Zizia aurea</i> , golden alexanders (early butterflies!)  | yel, 1- 3', self-sows; sun-lt shade; moist- wet |
| |  Grasses & Sedges  | |
| | <i>Agrostis hyemalis/scabra</i> , southern/northern tickle grass | small grass, airy fl stalks; dry-moist, pt shade-sun |
| | <i>Agrostis perennans</i> , Autumn bent-grass | grass, 15-40", delicate panicles; dry, sun |
| | <i>Andropogon gerardii</i> , big bluestem | tall showy grass; sun-pt sun; reg-wet |

NOTES – reg or "regular" is equivalent to typical garden soil, watered as needed.  = Good for pollinators
 Spring **Ephemeral**= E (foliage dies back by midsummer). * =Not strictly local native but regional/naturalized.

| | |
|--|--|
| <i>Carex eburnea</i> , bristleleaf sedge | 6-8" fine grass-like; shade-p shade, reg-dry, excellent drainage |
| <i>Carex granularis</i> , pale/limestone meadow sedge | glaucus leaves, corncob fruit, 2'; adaptable, limey, moist, sun-shade |
| <i>Carex grayi</i> , Asa Grey's sedge (scarce) | 2-3', large spiny seedhead; pt sun, reg soil |
| <i>Carex hystericina</i> , Porcupine sedge (scarce) | 2-3', large spiny seedhead; pt sun, reg soil |
| <i>Carex lurida</i> , sallow sedge | 2.5 ft; yellowish spiked seed heads; sun-pt sun; wet-moist |
| <i>Carex plantaginea</i> , plantain leaf/seersucker sedge | wide leaves, 6-12", evergreen; pt shade-shade, reg-moist |
| <i>Deschampsia cespitosa</i> , tufted hair grass | clumping grass, 2-3'; pt shade, reg-moist soil |
| <i>Dichanthelium clandestinum</i> , deertongue grass | erect grass, 2-3'; part shade-sun, moist-reg soil |
| <i>Elymus canadensis</i> , Canada wild rye | grass, blue-grn leaf, 2-3'; dry-wet, pt shade-sun |
| <i>Elymus hystrix</i> , bottlebrush grass | 2-3' grass; pt shade - shade, moist - dry |
| <i>Elymus riparius</i> , riverbank wild rye | 2-3' grass; pt sun-sun, wet-reg |
| <i>Elymus virginicus</i> , Virginia wild rye | 2-3' grass; sun-shade, wet-moist |
| <i>Panicum virgatum</i> , switch grass | 4' grass, wispy seed head; sun-p sun, dry-moist |
| <i>Schizachrium scoparium</i> , little bluestem (insect host) | grass, 3'sun, well-drained soil |
| <i>Tridens flavus</i> , purpletop tridens | grass, red- purp seedhead, 4', host plant; sun-p sun; reg-dry |
| Shrubs, Trees, Vines | |
| <i>Asimina triloba</i> , paw-paw (western NY) | small tree, purple-red fl, edible, colonizes; p shade-sun; moist |
| <i>Cephalanthus occidentalis</i> , buttonbush | Small shrub, white fl; Wet-moist, sun-pt sun |
| <i>Clematis virginiana</i> , virgin's bower | white fall flowering vine; moist, sun - part sun |
|  <i>Cornus alternifolia</i> , pagoda dogwood  | white fl, dark berries, small tree; pt shade - sun, reg- moist |
| <i>Cornus amomum</i> , silky dogwood | white fl, blue berries, shrub; p shade-sun, reg-moist |
|  <i>Cornus racemosa</i> , grey dogwood  | colony forming shrub, white fr; sun-pt shade, dry-wet |
| <i>Corylus americana</i> , American hazelnut | tall shrub, thickets, edible; dry-moist, sun-pt sun |
|  <i>Diervilla lonicera</i> , northern bush honeysuckle | low shrub (2'), stoloniferous, sm yellow flr; sun- pt shade, moist - dry |
|  <i>Hypericum prolificum</i> , shrubby St. John's wort | yellow, shrub to 5'; sun-pt shade, reg, well-drained soil |
| <i>Ilex verticillata</i> , winterberry | 3-12' shrub, bright fruit (female); sun-pt sun, wet-moist, pref acidic |
| <i>Juniperus virginiana</i> , eastern redcedar | Med evergreen tree, blue berries (female); sun, dry-reg soil |
| <i>Lindera benzoin</i> , spicebush  | yel/green fl, 6-12' shrub, spicy fragrance, yel fall; sun-pt sh, moist-dry |
|  <i>Liriodendron tulipifera</i> , tulip tree | very large tree, yel & orange fl; sun, moist, rich soil |
|  <i>Lonicera sempervirens</i> , coral honeysuckle | red/or/yel fl vine, 3-20'; sun-pt shade; rich moist (well drained!)soil |
| <i>Maclura pomifera</i> , Osage orange (naturalized, native AR,OK,TX) | large tree, green "brain" fruit on females; sun-pt shade, dry-reg |
| <i>Physocarpus occidentalis</i> , ninebark | shrub to 6', white fl; sun-pt shade, dry-moist |
| <i>Prunus serotina</i> , black cherry   | med large tree, edible fr; sun-pt sun, moist, well drained |
| <i>Prunus virginiana</i> , chokecherry | med tree, white fragrant fl, colonizes; sun, dry-moist, well drained |
| <i>Quercus sp</i> , oak – all wildlife! | med-tall tree, fall color; dry-moist, sun-pt sun |
| <i>Rhododendron prinophyllum</i> , early azalea (poisonous) | Pink, fragrant, 6-10' shrub, early; moist-wet; p shade-p sun |
| <i>Rhus aromatica</i> , fragrant sumac | 2-6' shrub, red fruit (female), fall color, suckers; sun-pt shade; adaptable |
| <i>Rosa carolina</i> , pasture rose | pink, 2-3', stoloniferous; sun-pt sh, dry-moist |
| <i>Rosa palustris</i> , swamp rose | pink, 2-6', stoloniferous;s un-pt sun, moist, acid-neutral soil |
| <i>Rubus odoratus</i> , purple-flowering raspberry | pink, shrub, edible fruit, spreads; dry-moist; sun- pt shade |
|  <i>Sambucus canadensis</i> , black elderberry | large shrub, white fl., edible black berry; sun-pt sun, reg soil |
|  <i>Sorbus decora</i> , northern mountain ash (garden source, poisonous) | med tree to 30'; white fl red berry; cooler sites, acidic moist, well drained, sun |
| <i>Spiraea alba</i> , meadowsweet | shrub to 6', white fl; sun, moist-wet, rich organic soil |
|  <i>Viburnum dentatum</i> , arrowwood (native NE US) | 6-10' shrub, white fl, tolerates black walnut; sun-pt shade; reg-moist, well drained |

NOTES – reg or "regular" is equivalent to typical garden soil, watered as needed.  = Good for pollinators
Spring **Ephemeral**= E (foliage dies back by midsummer). * =Not strictly local native but regional/naturalized.